

AD-A255 016



2

Final Report

Sources of Cloud Nuclei and Development of  
Remote Sensing Techniques for Cloud Parameters

Sean A. Twomey  
Principal Investigator

DTIC  
ELECTE  
AUG 27 1992  
S A D

Institute of Atmospheric Physics  
The University of Arizona  
Tucson, AZ 85721

June 1992

This document has been approved  
for public release and sale; its  
distribution is unlimited.

92 8 08

180800  
92-21215  
208

## Final Report

**Contract Title:** Sources of Cloud Nuclei and Development of Remote Sensing Techniques for Cloud Parameters

**Number:** N00014-90-J1288

**Principal Investigator:** S. Twomey, Institute of Atmospheric Physics,  
The University of Arizona

**ONR Program Manager:** Dr. W. A. Hoppel

The research focused on the problem of remotely determining the effective drop radius and optical thickness of marine stratus, and interpreting the results in terms of cloud susceptibility. Cloud susceptibility is here defined as the increase in albedo resulting from the addition of one cloud drop per cubic centimeter holding the cloud liquid water constant. The remote sensing technique employed made use of the AVHRR spectral channels on board the NOAA polar orbiting satellites. The method required an assumption of the cloud liquid water content which, for the marine stratus studied, was assumed to be  $0.3 \text{ g/m}^3$ .

As by-products of cloud susceptibility, cloud optical thickness and effective droplet radius was also determined. The values of these two parameters were compared to in situ measurements. Some discrepancies were noted, particularly in cloud optical thickness.

The retrieved range of cloud susceptibilities varied by about 2 orders of magnitude, from as low as  $0.23 \times 10^{-3}$  in stratus off the west coast of southern Africa to about  $20 \times 10^{-3}$  off the west coast of North America. California valley fog had values as low as  $0.05 \times 10^{-3}$ , extending the measured range for all clouds studied to almost three orders of magnitude. Studies indicated that ship tracks are less susceptible than non-track regions, due to the contamination of the tracks by CCN from the ships effluent. The susceptibility in and out of track may only differ by a factor of 2 to 4, or as high as 30 for their stratus.

Statement A per telecom Gary Geernaert  
ONE/Code 1122  
Arlington, VA 22217-5000

NWW 8/25/92

DTIC QUALITY INSPECTED 3

of magnitude, Africa to about fog had values studs studied to tracks are less tracks by CCN only differ by